



Research Article

New Distribution Record and Habitat Characteristics of Stinging Catfish (*Heteropneustes nani* Hossain, Sarker, Sharifuzzaman and Chowdhury, 2013) from Nepal

Asmit Subba^{1*}, Jash Hang Limbu² and Laxman Khanal^{1*}

¹Central Department of Zoology, Institute of Science and Technology, Tribhuvan University, Kathmandu 44618, Nepal

²College of Fisheries and Life Science, Shanghai Ocean University, Shanghai, China

Article History

Received: May 04, 2023

Revised: March 27, 2024

Accepted: March 27, 2024

Published: April 22, 2024

Authors' Contributions

AS and LK conceptualized the study. AS performed fieldwork, analyzed data and prepared the draft. JHL assisted in data analysis and revised the manuscript. LK supervised the study and finalized the manuscript.

Keywords

Catfish, Fish taxonomy, Jalthal Forest, Lowland Nepal, New records



Copyright 2024 by the authors. Licensee ResearchersLinks Ltd, England, UK. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract | The Stinging catfish (*Heteropneustes nani*) was described in 2013 from the Noakhali of Bangladesh through detailed morphological characterization. They differ significantly from their congeners based on small size, completely separated pectoral spine from rays bearing 9–10 serrae along the inner edge and united anal-caudal fin. In Nepal, only one species of *Heteropneustes*, i.e., *H. fossilis* has been reported. Based on the morphometric and meristic characteristics, this study reports the first-ever record of another species (*H. nani*) under the genus from eight locations in Jhapa District, eastern lowland Nepal, along with *H. fossilis* records. The two species distinctly differ, with *H. nani* conspicuously smaller than *H. fossilis*. *Heteropneustes nani* has been recorded from nutrient-rich, shallow, and turbid water bodies with abundant invasive aquatic macrophytes (*Eichhornia crassipes*, *Hydrilla verticillata*, and *Ceratophyllum submersum*). The type locality of the *H. nani* in Bangladesh and the newly reported locality in Nepal share similar tropical monsoon climates and river connectivity that might have facilitated their dispersal. Further studies are warranted to understand the detailed taxonomy and distribution pattern of *H. nani* in Nepal.

Novelty Statement | This study reported the occurrence of *Heteropneustes nani* from Nepal using field surveys from water bodies of eastern lowland of the country. Further, it characterized the habitats of the species and proposed the dispersal route from its original description locality in Bangladesh.

To cite this article: Subba, A., Limbu, J.H. and Khanal, L., 2024. New Distribution record and habitat characteristics of stinging Catfish (*Heteropneustes nani* Hossain, Sarker, Sharifuzzaman and Chowdhury, 2013) from Nepal. *Punjab Univ. J. Zool.*, 39(1): 15-20. <https://dx.doi.org/10.17582/journal.pujz/2024/39.1.15.20>

Introduction

Heteropneustes Müller, 1840 is a genus of catfish with stinging capabilities, known for its unique feature of having an air sac. This catfish is part of the Heteropneustidae

family in the order Siluriformes. *Heteropneustes* is found across a wide range of areas, including the Indian subcontinent and the mainland of several south and southeast Asian countries like India, Myanmar, Nepal, Pakistan, Sri Lanka, Vietnam, and Thailand (Arunachalam *et al.*, 1890; Berra, 2007; Ratmuangkhwang *et al.*, 2014). They inhabit ditches, swamps, canals, marshes, ponds, and waterlogged areas of lowlands with shallow water depth and low dissolved oxygen levels. They are characterized by

Corresponding Author:

Asmit Subba, subbaasmit926@gmail.com;

Laxman Khanal, laxman.khanal@cdz.tu.edu.np

an elongated and compressed body with short dorsal and long pelvic fins, a pectoral serrated spine associated with venom glands and four pairs of barbels, a pair of elongated and tubular cavities of air sacs extend posteriorly almost up to the tail, and adaptation to oxygen-deficient waters (Nelson, 2006; Berra, 2007).

Five nominal species of *Heteropneustes* are described, namely *H. microps* (Pethiyagoda and Bahir, 1998) and *H. longipetrolis* (Devi and Raghunathan, 1999) from Western Ghats, India; *H. kemratensis* (Fowler, 1937) from the Chao Praya River in Thailand; *H. fossilis* (Bloch, 1794) from Tamil Nadu, India; and *H. nani* from Noakhali, Bangladesh (Hossain *et al.*, 2013). *Heteropneustes nani* stands out from its relatives due to its notably smaller size, with an average length of 98.08 mm from the snout to the tail (compared to 223.96 mm in *H. fossilis*, 152.40 mm in *H. microps*, 210.00 mm in *H. kemratensis*, and 150.00 mm in *H. longipetrolis*). Additionally, they have a unique combination of features: Their head is larger compared to their body depth; they have small jaws with fleshy lips and a terminal mouth; their body is dark colored with faint lateral bands on the sides; and their pectoral spines are set apart from the rays featuring 9–10 serrations along the inner edge (Hossain *et al.*, 2013).

The type locality of *H. nani*, the Noakhali area of Bangladesh, lies adjacent to Nepal, where the species has not been reported despite the resemblance of suitable habitats (lakes, ponds, marshes, and floodplains) between eastern lowlands of Nepal and Bangladesh. Only a single species of the genus, *H. fossilis*, was reported from Nepal (Shrestha, 2008). Morphologically, *H. fossilis* differs from *H. nani* in multiple characters. The upper body of *H. fossilis* is either yellow or dark purplish-brown color, while the lower body is comparatively lighter. The mouth extends backward, and the pectoral spine is connected to the fin rays by a membrane featuring 3–4 serrations that curve forward along the inner edge at the front tip. Additionally, there is a clear separation between the anal and caudal fins, marked by a distinct notch, unlike in *H. nani* (Bloch, 1794; Shrestha, 2008; Hossain *et al.*, 2013). *Heteropneustes fossilis* live in freshwater environments, occasionally in slightly salty or brackish water such as ponds, ditches, swamps, marshes, and muddy rivers. During the dry season, they can survive in partially wet and semi-dry mud, as noted by De Silva *et al.* (2015) and Fernando and Goonatilake (2019). It is one of the common species distributed along the lowland and midland waters of Nepal (Shrestha, 2008).

As a result of its altitudinal variation and geographical location intertwining the eastern and western Himalayas, Nepal is home to a wide variety of fish species (Shrestha, 2000). However, due to limited studies, many wildlife species, including fish, remain unidentified and lack proper taxonomic descriptions (Limbu *et al.*, 2023).

Comprehensive research involving both meticulous classification efforts and genetic examinations is essential to formulate effective strategies for inventory and conserving the country's fish population (Khatri *et al.*, 2020). This study marks the first record of *Heteropneustes nani* in Nepal from the eastern lowlands of Jhapa District. The habitat characteristics of the newly reported *H. nani* in Nepal have also been explained.

Materials and Methods

On 5th September 2021, during the Jalthal Biodiversity Profiling Survey, the first author observed a specimen from the small pond near the Deuniya River bridge, Jhapa District, Koshi Province, Nepal (Figure 1) where local fishermen had collected the fish. In a close observation, it appeared remarkably different from the previously reported stinging catfish from the locality, i.e., *H. fossilis*. Morphologically, they were smaller than *H. fossilis* and had completely separated pectoral spine from rays, and had 9–10 serrae along the inner edge, similar to the explanation of Hossain *et al.* (2013) for *H. nani*. Further explorations were done, and similar specimens were collected from eight different locations in Jhapa District between September 2021 and December 2022. Additional information, including habitat, local species names, locations, season, and abundance, was also collected.




Figure 1: A map representing the locations from where *Heteropneustes nani* samples were collected. Each colour symbolizes the respective land use polygons.

The specimens were collected from small lakes, ponds, marshes, and flood plains of Deuniya River (88.050731°E, 26.530253°N), Kankai River (87.838364°E, 26.541983°N), Kamal River (87.792928°E, 26.554314°N), Biring River (87.900111°E, 26.537231°N), Thade Pokhari (87.945086°E, 26.451831°N), Kechana Jheel areas

(87.999803°E, 26.383389°N), Jalthal Forest wetlands (87.986953°E, 26.454278°N), and Tindhwang areas (87.798356°E, 26.588661°N) with the help of local fishermen. These areas had abundant invasive aquatic macrophytes (*Eichhornia crassipes*, *Hydrilla verticillata*, and *Ceratophyllum submersum*), with low water depth and high turbidity, typical of nutrient-rich waters with high eutrophication rates that lead to low dissolved oxygen. In each station, a single specimen was collected.

The collected specimens were photographed and preserved in 95% alcohol in a plastic jar by making their caudal part upright to protect their caudal fin for further analysis. To make sample tracking easier, a collection number was given to each sample. The collected samples (*H. fossilis* and *H. nani*) were brought to the Central Department of Zoology, Tribhuvan University, for detailed taxonomic identification and comparison. The specimens are deposited in the Museum of the Central Department of Zoology, Tribhuvan University (CDZMTU-HN1). Since only two individuals of *H. fossilis* were collected, we could not use the individual data for morphometric comparison. Instead, we used adult specimen data collected by Hossain *et al.* (2013).

A diagnostic morphometric measurement was obtained based on Hubbs and Lagler (2004), which includes: Snout-vent length (SVL or SL); head length (HL); head width (HW); snout length (SNL); eye diameter (ED); body depth (BD); anal fin length (AFL); dorsal fin length (DFL); pectoral fin length (PFL); pectoral spine length (PSL); caudal fin length (CFL); first barbel (BL1); second barbel length (BL2); third barbel length (BL3) and fourth barbel length (BL4). The measurements were taken using a digital Vernier Caliper to the nearest tenth of a millimeter from the specimen's left side following the method of Ng and Wright (2009) and were presented as mean±SD (standard deviation). The collected information was tabulated and visualized using unsupervised cluster analysis.

Results and Discussion

Based on the detailed diagnostics and morphometric data, the current species was smaller (95.7–98.97) mm, dorsally blackish, and lighter in ventral color in live specimens, and short and sub-cylindrical up to the pelvis base (Figure 2). Their head is generally depressed with length (14.11–17.21 with a mean of 15.47% of SL) less than the body depth (13.80–15.67 with a mean of 14.78714% of SL). They have a terminal mouth, villiform bands on jaws, small eye diameter (8.87–12.33 with a mean of 10.819 % HL), and four pairs of barbels; 34.23–36.54, 27.43–29.46, 30.28–31.23, and 35.11–36.42% of HL in order, respectively.

Heteropneustes nani differs from the congener *H. fossilis*, which had already been documented from Nepal,

in having separated pectoral spines and nine to ten antrorse serrae, whereas the latter has pectoral spines attached by a membrane. Similarly, *H. nani* has a united caudal fin, while *H. fossilis* has a distinct caudal notch (Hossain *et al.*, 2013). In addition, other morphological features distinctively separate *H. nani* and *H. fossilis* (Table 1 and Figure 3). The unsupervised cluster analysis of morphological characters revealed two distinct clusters, *H. nani* and *H. fossilis*, with *H. fossilis* plotting on the positive side of the first principal component (Figure 4), reflecting the morphological distinctness between the two species, and the conspicuously smaller size of *H. nani*.



Figure 2: Photographs of *Heteropneustes nani* recorded from the study area; Dorsal (A) and Ventral (B) view.



Figure 3: The illustration depicting two species of *Heteropneustes* found in Nepal. Plate A shows a sub-adult *Heteropneustes fossilis*, while plate B features a *Heteropneustes nani*.




Figure 4: Cluster-wise representation of *H. fossilis* and *H. nani* into two separate clusters, one of each species.

Table 1: Representing summary of diagnostic morphometric characteristics between *H. nani* and its congeners *H. fossilis*.

Features	<i>Heteropnuestes nani</i> (n= 8)		<i>Heteropnuestes fossilis</i> (n=12)*	
	Range	Mean±SD	Range	Mean±SD
Head length	14.11–17.21	15.47±1.183	12.77–14.56	13.65±0.65
Body depth	13.80–15.67	14.78±1.29	13.10–15.50	14.46±0.79
Anal fin length	61.88–66.30	63.44±1.827	59.47–63.11	60.89±1.27
Dorsal fin length	12.24–13.61	12.95±0.519	9.09–10.12	9.54±0.46
Pectoral fin length	10.87–12.35	11.73±0.533	10.07–10.55	10.25±0.16
Pectoral spine length	8.86–9.63	9.28±0.25	7.78–9.03	8.37–0.48
Caudal fin length	8.56–9.83	8.995±0.460	8.33–10.72	9.82±0.98
Ventral fin length	7.43–8.07	7.781±0.23	7.69–8.40	7.93±0.24
First barbel length	34.23–36.54	36.23±0.83	8.71–21.96	17.32±4.77
Second barbel length	27.43–29.46	28.83± 0.71	22.25–31.98	28.44±3.33
Third barbel length	30.28–31.23	30.83±0.33	19.30–27.71	23.68±2.73
Fourth barbel length	35.11–36.42	35.77±0.4	18.32–22.69	20.57±1.89
Eye diameter	8.87–12.33	10.819±1.12	11.24–12.18	11.71±0.47
Head width	88.20–94.46	91.69±2.226		
Pectoral spine	Completely separated from rays		Attached to rays by a membrane	
Anal–caudal fin	United		Separated by a distinct notch	

*Source: (Hossain *et al.*, 2013).

Our study area lies in a tropical monsoon climate zone with adequate rainfall, rich in agricultural (paddy) landscape with several ponds, ditches, swamps, rivers, floodplains, and small lakes with a temperature range from 10.10°C to 33.35°C respectively (Bhattarai, 2017; Bhattarai *et al.*, 2017). The climatic conditions of the *H. nani* localities in Nepal and Bangladesh are nearly similar, i.e., strong effects of the south-Asian summer monsoon (Islam, 2009; Kamruzzaman *et al.*, 2018). Besides climatic similarity, the riverine network connectivity has been found to affect the pattern of fish biodiversity because fish have to follow watercourses between habitats (Lynch *et al.*, 2013; Shao *et al.*, 2019). There are river connections between the localities of the original description of the *H. nani* from Bangladesh and the location of the present records in Nepal (Figure 5), which provided dispersal pathways.

The Kankai River and its tributaries in Nepal flow southward, transitioning into the Mahananda River and then the Ganges River as they cross into India. Eventually, these waters merge into the Padma River, which further flows into the Bay of Bengal through Bangladesh (Azam *et al.*, 2018; Dewan *et al.*, 2017). This connectivity of rivers might have facilitated the dispersal of *H. nani* across the river network. Further studies are required to document the fine-scale distribution of the species across water bodies in Nepal and India.



Figure 5: Map depicting the river connections between the original description of *H. nani* in Bangladesh (Hossain *et al.*, 2013) and present records of the species in Nepal.

The low oxygen and nutrient-rich lentic ecosystem favors tubular air cavities containing *Heteropnuestes* species (Berra, 2007). The current study shows that *H. nani* distribution is widespread across the lowlands of Jhapa District together with *H. fossilis*. Through interviews, we found the local names of *H. nani* and *H. fossilis* are Kachad and Singhi, respectively, based on their pectoral poisonous spines. *H. nani* was recorded from more localities compared to *H. fossilis*, likely because the surveys were conducted in habitats that were more preferable for *H. nani*. Also, we

recorded *H. fossilis* from comparatively deeper water levels than *H. nani* which were mostly in a purer waterside area.

The observed occurrence of the species can be explained by environmental variables such as temperature, precipitation, and habitat vegetation of our study area which resemble those of Noakhali in Bangladesh, from where the species was first recorded. In addition, other congeners, i.e., *H. microps*, *H. longipetrolis*, and *H. kemratensis*, are geographically distant from the survey locality and the morphological characteristics are closer to those of *H. nani*, compounding supporting that the specimens we collected from the eastern lowland of Nepal belong to *Heteropneustes nani*.

Conclusion and Recommendations

This research has verified the presence of *Heteropneustes nani* in the eastern lowlands of Nepal. Despite coexisting with similar species like *H. fossilis*, they exhibit notable differences in morphology and habitat characteristics. These distinctions include the size, shape, coloration, the count of anterior serrae, and the separation of pectoral spines. Moreover, *H. nani* is widely distributed in shallow water bodies across the Jhapa District, showing a preference for oxygen-scarce lentic water ecosystems compared to *H. fossilis*. Furthermore, we assume that *H. fossilis* found on comparatively deeper, less turbid ditches and wetlands loaded with agricultural chemical-rich water than *H. nani*.

This study revealed a direct connection to the type locality of *H. nani* based on their river network linkage, although we need concrete evidence regarding their migration patterns. Based on our hypothesis, it is plausible that *H. nani* may be distributed across the southern areas of West Bengal, Bihar, and Bangladesh, as these regions feature similar wetland ecosystems with comparable physiochemical and environmental characteristics. Further studies, employing both morphological and genetic data to comprehensively confirm the taxonomic status of the species, are needed. Such investigations would also contribute to a better understanding of the distribution of *H. nani* throughout Nepal.

Acknowledgements

We acknowledge financial support from Forest Action Nepal under the Darwin Initiative Project UK funded project (Ref. 26-022), and all the members who participated actively for the collection of species. We also thank the Museum of the Central Department of Zoology for storing the specimens.

Conflict of interest

The authors have declared no conflict of interest.

References

- Arunachalam, M., Johnson, J.A. and Manimekalan, A., 1890. New record of *Heteropneustes microps* (Gunther) (Claridae: Heteropneustidae) from Western Ghats River, India. *J. Bombay Nat. Hist. Soc.*, **68**: 1–492. <https://www.biodiversitylibrary.org/item/18553>
- Azam, M.M., Kumari, M., Maharana, C., Singh, A.K. and Tripathi, J.K., 2018. Recent insights into the dissolved and particulate fluxes from the Himalayan tributaries to the Ganga River. *Environ. Earth Sci.*, **77**: 313. <https://doi.org/10.1007/s12665-018-7490-7>
- Berra, T.M., 2007. *Freshwater fish distribution*. The University of Chicago Press: Chicago, USA. <https://doi.org/10.7208/chicago/9780226044439.001.0001>
- Bhattarai, K.P., 2017. Enumeration of flowering plants in Tarai Sal (*Shorea robusta* Gaertn.) Forest of Jalthal, Eastern Nepal. *J. Plant Resour.*, **15**: 14–20. <https://doi.org/10.3126/on.v14i1.16438>
- Bhattarai, K.P., Shrestha, P.K. and Koirala, O., 2017. Physico-chemical properties of water and soil of Bhimsen Pokhari Wetland, Jhapa District, Eastern Nepal. *Nepal J. Biosci.*, **7**: 72–76. <https://doi.org/10.3126/njbs.v7i1.41785>
- Bloch, M.E., 1794. *Ichthyologie, ou Histoire naturelle, generale et particulier des poissons*. A Berlin, Chez l'auteur, and chez François de la Garde libraire.
- De Silva, M., Hapuarachchi, N., Jayaratne, T., Robeson, D., and Wildlife Conservation Society, G., 2015. *Sri Lankan freshwater fishes*. Wildlife Conservation Society, Galle.
- Devi, K.E. and Raghunathan, M.B., 1999. *Heteropneustes longipetrolis* (siluriformes: heteropneustidae) a new species from the Anamalai Hills, in the Western Ghats. *Rec. Zool. Surv. Ind.*, **97**: 109–115. <https://doi.org/10.26515/rzsi/v97/i3/1999/160129>
- Dewan, A., Corner, R., Saleem, A., Rahman, M.M., Haider, M.R., Rahman, M.M., and Sarker, M.H., 2017. Assessing channel changes of the Ganges-Padma River system in Bangladesh using Landsat and hydrological data. *Geomorphology*, **276**: 257–279. <https://doi.org/10.1016/j.geomorph.2016.10.017>
- Fernado, A. and Goonatilake, D.A., 2019. *Heteropneustes fossilis*. IUCN red list of threatened species 2019: e.T166452A60585129. <https://dx.doi.org/10.2305/IUCN.UK.2019-3.RLTS.T166452A60585129.en>. Accessed on 02 September 2023.
- Fowler, H.W., 1937. Zoological results of the third de Schauensee Siamese expedition. Part VIII: Fishes Obtained in 1936. *Proc. Acad. Nat. Sci. Phila.*, **89**: 125–264.
- Hossain, M.S., Sarker, S., Sharifuzzaman, S.M. and Chowdhury, S.R., 2013. New species of stinging catfish *Heteropneustes nani* (Siluriformes:

- Heteropneustidae) from Noakhali, Bangladesh. *Vertebr. Zool.*, **63**: 259–267. <https://doi.org/10.3897/vz.63.e31441>
- Hubbs, C.L. and Lagler, K.F., 2004. *Fishes of the great lakes region, revised edition*. University of Michigan Press. <https://doi.org/10.3998/mpub.17658>
- Islam, M.N., 2009. Rainfall and temperature scenario for Bangladesh. *Open Atmospheric Sci. J.*, **3**: 93–103. <https://doi.org/10.2174/1874282300903010093>
- Kamruzzaman, M., Rahman, A.T.M.S., Ahmed, M.S., Kabir, M.E., Mazumder, Q.H., Rahman, M.S. and Jahan, C.S., 2018. Spatio-temporal analysis of climatic variables in the western part of Bangladesh. *Environ. Dev. Sustain.*, **20**: 89–108. <https://doi.org/10.1007/s10668-016-9872-x>
- Khatri, K., Jha, B.R., Gurung, S. and Khadka, U.R., 2020. Freshwater fish diversity and its conservation status in different water bodies of Nepal. *Nepal J. Environ. Sci.*, **8**: 39–52. <https://doi.org/10.3126/njes.v8i1.34442>
- Limbu, J.H., Rajbanshi, D., Subba, B.R., Subba, A., Yang, J.Q. and Li, C., 2023. First record of catfish *Amblyceps waikhomi* (Darshan, Kachari, Dutta, Ganguly, and das 2016) (Siluriformes: Amblycipitidae) for Nepal from the Singhiya River of Morang District, Eastern Nepal. *Int. J. Zool.*, **2023**: 3707208. <https://doi.org/10.1155/2023/3707208>
- Lynch, H.J., Campbell, G.E.H., Muneeppeerakul, R., Arunachalam, M., Rodriguez-Iturbe, I. and Fagan, W.F., 2011. How restructuring river connectivity changes freshwater fish biodiversity and biogeography. *Water Resour. Res.*, **47**: W05531. <https://doi.org/10.1029/2010WR010330>
- Nelson, J.S., 2006. *Fishes of the World* (4th ed.). John Wiley and Sons, Inc., Hoboken, New Jersey. pp. 707.
- Ng, H.H. and Wright, J.J., 2009. A new torrent catfish from western Thailand (Siluriformes: Amblycipitidae). *Copeia*, **2009**: 369–377. <https://doi.org/10.1643/CI-08-113>
- Pethiyagoda, R. and Bahir, M.M., 1998. *Heteropneustes microps*, a junior synonym of *H. fossilis* (Osteichthyes: Heteropneustidae). *J. South Asian Nat. Hist.*, **3**: 113–114.
- Ratmuangkhwang, S., Musikasinthorn, P. and Kumazawa, Y., 2014. Molecular phylogeny and biogeography of air sac catfishes of the *Heteropneustes fossilis* species complex (Siluriformes: Heteropneustidae). *Mol. Phylogenet. Evol.*, **79**: 82–91. <https://doi.org/10.1016/j.ympev.2014.05.009>
- Shao, X., Fang, Y., Jawitz, J.W., Yan, J. and Cui, B., 2019. River network connectivity and fish diversity. *Sci. Total Environ.*, **689**: 21–30. <https://doi.org/10.1016/j.scitotenv.2019.06.340>
- Shrestha, T.K., 2008. *Ichthyology of Nepal: A study of fishes of the Himalayan waters*. pp. 388.
- Shrestha, T.K., 2000. Cold water fisheries development in Nepal. In: *Cold water fisheries in the Trans-Himalayan countries* (eds. T. Petr and D.B. Swar, D.B.). pp. 47–58.

